


Inelastic UV Scattering as a new Technique to Investigate Collective Excitations in Condensed Matter Physics

C. Masciovecchio

Elettra Synchrotron, S.S. 14 km 163,5 – 34012 Basovizza, Trieste – Italy

- **Inelastic UV Scattering (IUVS) with μeV Energy Resolution**
- **The Beam-line Design and Construction**
- **Applications to Condensed Matter Physics and Preliminary Results**
- **Conclusions**

Inelastic UV Scattering with μeV Energy Resolution






$$I_{if}(\mathbf{Q}, \omega) = \frac{I_0 k_f^4}{R} \int_{-\infty}^{\infty} dt e^{-i\omega t} \langle \delta\alpha_{if}(\mathbf{Q}, 0) \delta\alpha_{if}(\mathbf{Q}, t) \rangle$$

Spectral Density of
Polarizability Tensor Fluctuations

Spherical Molecules: $I_{VV}(\mathbf{Q}, \omega) = \alpha^2 S(\mathbf{Q}, \omega)$

Density Fluctuation
Spectrum

Available probes to investigate collective excitations



Investigations in the Intermediate Region could Shed Light on:

Liquids - Fluids

- Transition from the **Hydrodynamic to the Kinetic** regime in Simple liquids and fluids.
- Effect of the **Local Structure** on the Collective Dynamics in Molecular liquids, Associated liquids, and H-bonded liquids with a specific interest in Water and Water Solutions.

Glasses

- Nature of the **Vibrational Modes** in the Mesoscopic space-time region.
- **Relaxational Processes** in Super-Cooled liquids and their relation to the Glass Transition.
- Vibrational and Relaxational **Low Temperature Properties** of Fragile and Strong glasses.

Resonant Scattering (Tunability**)**

- Low **count-rate** experiments.
- Determination of **Partial Dynamic Structure Factor**.
- **Transverse Dynamics** of the system.

Experimental requirements for IUVS

- Incident **Energy** in the 5 – 11(30) eV range ($\lambda \approx 240 - 110(40)\text{nm}$)
- High incident photon **Flux** on the Sample ($> 10^{12} \text{ photons/s}$)
 \swarrow $\sim 1 \text{ counts/s}$
- High **Resolving Power** ($\approx 10^5 - 10^6$)

$$\omega \approx 10 - 10^3 \mu\text{eV}$$

$$Q \approx 0.02 - 0.4 \text{ nm}^{-1}$$

The Beamline Design and Construction


Linear Undulator ?

4.5 m length, 125 mm period, 400 mA

$2 \cdot 10^{15}$ photons/s/0.1% bandwidth

1.5 kW on the first mirror

Figure 8 solution



$$N_p = 32$$


$$\lambda = 140 \text{ mm}$$

$$K_x = 3.4$$

$$K_y = 9.4$$


22 W on first mirror !!

$2 \cdot 10^{15}$ photons/s/0.1% BW ($2 \cdot 10^{12}$ photons/s)



The NIM Monochromator

Normal Incidence Monochromator **NIM** (*Czerny-Turner* design)



Monochromator & Analyzer design

$$\frac{\Delta E}{E} = \frac{\delta \cdot \text{ctg} \theta}{2F} = \frac{50 \mu\text{m} \cdot \text{ctg}(70^\circ)}{16\text{m}} \approx 1 \cdot 10^{-6}$$

The Beamline

Scanning Resolution: 80 *nrad*

Autocollimator Control: 50 *nrad*


Slits Opening: 5 – 12000 μm

Grating Dimensions 420 × 220 *mm*


Monochromator Length 8 *m*

Vacuum: 10^{-8} *mbar* (stability)


Temperature Variation: 0.01°C



The Construction



The Slits




Four independent Blades design

Resolution = parallelism = repeatability : $1 \mu\text{m}$

Rotation range (resolution): 4 mrad ($2 \mu\text{rad}$)

Translation range (resolution): 50 mm ($5 \mu\text{m}$)

The Gratings




Efficiency: 60% @ 250 nm
50% @ 200 nm
Ghost Intensity $< 1 \cdot 10^{-4}$ PL
Slope Errors ~ 0.2 arcsec

Dimensions: 420×220 mm
Groove Density: 52.67 g/mm
Coating: Al + MgF₂ (25 nm)
Blank Material: Zerodur

The Scanning System

Renishaw Head: RGH 25 UHV
Analogue Reader: RGB 25 B
Sinusoidal Interpolator: MXH 250
(20 nm resolution – 80 nrad)

Nanomotion HR8 UHV Piezo Motor
Minimum movement: 1 nm




Monochromator + Analyzer



Sample Stage



Sample Manipulator




Nanomotion Motors to
achieve 1 μ rad resolution
movement

Detectors

1) **CCD** Thinned Back Illuminated

Chip EEV 4210 13,5 μm pixel size (2048×515)

Peltier cooled @ -90°C  1 *el/pixel/hour*



2) **MCP** coupled to a Resistive Anode

30 μm spatial resolution


Single Photon Counter



3) **Photomultiplier**


no spatial resolution

Single Photon Counter



The Resolution

Resolution measured using copper scattering



$$\Delta E \approx 2 \cdot 10^{-6}$$

$$E \approx 5 - 11 \text{ eV}$$

$$\text{Flux} \approx 10^{12} \text{ ph/s}$$

Applications to Condensed Matter Physics and Preliminary Results

Glass-Forming Systems


Puzzling properties

Glass-transition mechanism

Relaxation processes

Thermal anomalies

Excess in the vibrational DOS



Knowledge of C_0 , C_∞ , $\tau_R(Q)$, $C(Q,T)$



Formulation of Models describing the
Glass Transition

Structural Relaxation in Water

First evidence of glassy water: **LDA** (*Burton et al.*, 1936) $T_g \approx 130 \text{ K}$

Water exhibits very unusual properties:

- Negative volume of melting
- Density maximum in the normal liquid range
- Isothermal compressibility minimum in the normal liquid range
- Increasing liquid fluidity with increasing pressure

Intermolecular forces and

Hydrogen Bond

($T \sim 4^\circ \text{C}$)

Evidence of a relaxation with a characteristic time of $\tau \sim 1 \text{ ps}$

high frequency investigations: **IXS** (*Sette et al.*, PRL 1996, *Monaco et al.*, PRE 1999)

low frequency investigations: **Ultrasonic** (*Slie et al.*, JCP1966), **BLS** (*Cunsolo et al.*, JCP1996)


Most Sensible Region $\omega_P \tau_R \sim 1$ $\tau_R (4^\circ \text{C}) \sim 3 \text{ ps}$ $\omega_R \sim 200 \text{ } \mu\text{eV}$

$$\omega_P = v \cdot Q = v \cdot n \cdot 4 \pi / \lambda \cdot \sin (\theta / 2) \sim 100 - 250 \text{ } \mu\text{eV}$$

IUVS first measurements

Water from liquid to undercooled state

- Cell: Fused Silica Fluorescence standard Cell
- Momentum Transfer: 0.9 nm^{-1} ($\lambda = 180 \text{ nm}$)
- Temperature range: $-12 \rightarrow 76 \text{ }^{\circ}\text{C}$



Conclusions

IUVS design started on March 2000. Today the beamline is in its commissioning phase and preliminary results show that it is now possible to perform inelastic scattering experiment in a Momentum-Energy transfer region not *accessible* by other experimental techniques.

The possibility to study disordered systems in the mesoscopic region will shed light on several open problems in the physics of these intriguing systems.

Acknowledgments

F. Sette, ESRF - France

G. Ruocco, University “La Sapienza” Rome - Italy

D. Cocco, A. Gessini, S. Di Fonzo, ID group, Detector Group, ELETTRA - Italy

S. Santucci, D. Fioretto, University of Perugia - Italy